

www.biolmedonline.com

## Hyperglycemia induced by subchronic co-administration of chlorpyrifos and lead in Wistar rats: Role of pancreatic lipoperoxidation and alleviating effect of vitamin C

<sup>1</sup>Ambali SF\*, <sup>1</sup>Shuaib K, <sup>1</sup>Edeh R, <sup>1</sup>Orieji BC, <sup>1</sup>Shittu M, <sup>2</sup>Akande M

<sup>1</sup>Toxicology Unit, Department of Veterinary Physiology and Pharmacology, Ahmadu Bello University, Zaria, Nigeria.

<sup>2</sup>Department of Veterinary Pharmacology and Toxicology, University of Abuja, Nigeria.

\*Corresponding Author: fambali2001@yahoo.com, atunluse@gmail.com

### Abstract

Studies were conducted to evaluate the role of pancreatic lipoperoxidation on hyperglycemia induced by subchronic co-administration of chlorpyrifos (CPF) and lead (Pb) in Wistar rats and the ameliorative effect of vitamin C. Forty male Wistar rats divided into 4 groups of 5 animals in each group were used for this study. Rats in group I were dosed with corn oil (2 ml/kg) while those in group II were dosed with vitamin C (100 mg/kg). Group III were co-administered CPF (4.25 mg/kg~ 1/20<sup>th</sup> LD<sub>50</sub>) and Pb (250 mg/kg~1/20<sup>th</sup> LD<sub>50</sub>) while those in group IV were pretreated with vitamin C (100 mg/kg) and then co-administered with CPF (4.25 mg/kg) and Pb (250 mg/kg) 30 min later. The regimen were administered once daily by gavage for a period of 9 weeks. The rats were sacrificed and serum obtained from the blood samples were analyzed for glucose concentration. The liver and pancreas samples were analyzed for glycogen and malonaldehyde (MDA) concentrations, respectively. The study showed that co-administration of CPF and lead caused increased glucose and MDA concentrations, and a reduced glycogen concentration. Pretreatment with vitamin C restored the concentrations of glucose, glycogen and MDA to apparently normal level. In conclusion, pretreatment with vitamin C restored the hyperglycemia and reduced glycogen concentration induced by co-administration of CPF and Pb partly due to its antioxidant properties.

**Keywords:** Chlorpyrifos; lead; hyperglycemia; lipoperoxidation; amelioration; vitamin C.

### Introduction

Man and animals are exposed to a mixture of chemical contaminants in the environment, which directly or indirectly affect their health and well-being. The environment is pervasive with multiple chemicals that directly or indirectly interact with each other and the ecosystem. Few studies have evaluated the effect of multiple chemical contaminants on human and animal health as efforts have centered on evaluating the effect of a single contaminant. Pesticides and heavy metals constitute the most widespread environmental contaminants due to their ubiquitous use in all aspects of human endeavor.

Organophosphate (OP) insecticides are one of the most widely used insecticides accounting for 50% of the global insecticidal use (Casida and Quistad, 2004). The compelling needs to improve human and animal nutrition and promote public health has led to an increase in OP usage in recent time as they are used extensively to control agricultural, household and structural pests (Pope, 1999). Although, neurotoxicity is the hallmark of OP insecticide poisoning, other systemic toxicity have been observed following acute or repeated exposure.

Hyperglycemia is one of the side effects of OP poisoning in humans with blood glucose rising by about five folds (Namba *et al.*, 1971; Hayes *et al.*, 1978; Meller *et al.*, 1981). Similarly, an epidemiological study has found a direct relationship between consistent exposure to OP insecticides (including CPF) and incidence of diabetes among the pesticide applicators (Montgomery *et al.*, 2008). Hyperglycemia has been observed following OP exposure in animal models (Seifert, 2001; Ambali, 2009).

Chlorpyrifos (CPF) is one of the most widely used OP insecticides in agriculture and public health. Due to their wide availability, poisoning by CPF is common (Garcia *et al.*, 2003) as residual amounts have been detected in the soil, water bodies, vegetables, grains and other food products (Poet *et al.*, 2004). Similarly, lead is one of the most pervasive heavy metal contaminants in the environment (Krishna and Ramachandran, 2009). Exposure to lead has been known to adversely affect human and animal health in urbanized communities (Wang *et al.*, 2006). Hyperglycemia is one of the signs associated with lead poisoning (Stevenson *et al.*, 1976; Shaffi, 1979).

Although many mechanisms are involved in both CPF and Pb poisoning, the induction of oxidative stress is central to the two contaminants (Ercal *et al.*, 1996; Gultekin *et al.*, 2001; Olaleye *et al.*, 2007; Ambali *et al.*, 2010a-d). The animal body has evolved an effective antioxidant system to combat the menace posed by oxidative stress. Vitamin C is one of the most important free radical scavengers in extracellular fluid, trapping radicals in the aqueous phase, and protecting biomembranes from peroxidative damage (Yavuz *et al.*, 2004). Vitamin C has shown promise in alleviating toxicity induced by CPF (Ambali *et al.*, 2007; 2010c,d) and lead (Houston and Johnson, 2000; Oladipo, 2010). It is therefore conceivable that pretreatment with vitamin C will reduce lipoperoxidative changes induced by co-administration of CPF and Pb. The aim of the present study is therefore to evaluate the alleviating effect of vitamin C on hyperglycemia and hepatic glycogen depletion induced by subchronic co-administration of CPF and Pb in Wistar rats.

## Materials and Methods

### Experimental animals

Forty 6-week old adult male Wistar rats were obtained from the Animal House of the Department of Veterinary Physiology and Pharmacology, Ahmadu Bello University, Zaria, Nigeria. The rats were fed on standard rat pellets and water provided *ad libitum*. The experimental procedures were conducted in accordance with the guideline on the use of laboratory animals (NRC, 1996).

### Chemicals

Commercial grade CPF (Termicot® 20% EC, Sabero Organics, Gujarat, India) was dissolved in soya oil to make 10% stock solution, which was subsequently used for the experiment. Analytical grade lead acetate (Kiran Light Laboratories, Mumbai, India) used for the study was made into a 20% stock solution in distilled water. Commercial grade vitamin C tablets (Emzor Pharmaceutical Ltd, Nigeria, BN: 618N) was prepared in distilled water to make 10% stock solution.

### Animal treatments

Forty weaned male Wistar rats were divided into four groups of 10 animals per group. The rats in group I were administered corn oil (2ml/kg), while those in group II were administered

Vitamin C (100mg/kg). Rats in group III were co-administered CPF [4.25mg/kg, ~1/20<sup>th</sup> LD<sub>50</sub> (Ambali, 2009)] and lead acetate [225mg/kg, ~1/20<sup>th</sup> LD<sub>50</sub> (Oladipo, 2010)], respectively. Rats in group IV were pretreated with vitamin C, and then co-administered with CPF (4.25mg/kg) and Pb (225mg/kg), 30 min later. These regimens were administered orally by gavage once a day for a period of 9 weeks. At the end of the study period, the rats were sacrificed by jugular venisection after light chloroform anesthesia. Serum obtained from each blood sample was used to evaluate the concentration of glucose while the liver was assayed for glycogen concentration. The pancreas was assayed for the concentration of malonaldehyde (MDA) as an index of lipoperoxidation.

### Evaluation of serum glucose concentration

The serum glucose concentration was determined using the glucose oxidase method. The principle of the method is based on the ability of glucose oxidase to catalyse the oxidation of β-D-glucose to D-glucono-σ-lactone with the concurrent release of hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>). In the presence of peroxidase (POD) this H<sub>2</sub>O<sub>2</sub> enters into a second reaction involving *p*-hydroxybenzoic acid and 4-aminoantipyrine with the quantitative formation of a quinoneimine dye complex which is measured at 510 nm.

### Evaluation of hepatic glycogen concentration

The hepatic glycogen concentration was evaluated using the gravitational method of Good *et al.* (1933). Briefly, 0.5g of liver was extracted with 3mls of 30% KOH, incubated for 30 min. at 100°C, and then brought to acidic pH by addition of 20% trichloroacetic acid. Precipitated protein was removed by centrifugation for 10 min at 3000xg. Glycogen was precipitated by ethanol and weighed. The results were expressed in g of glycogen/100g of liver sample.

### Evaluation of pancreatic lipoperoxidation

The MDA concentration of the pancreas was assayed using the double heating method of Draper and Hadley (1990). 0.3g of pancreas was homogenized in 30 ml cold phosphate buffered saline to obtain and centrifuged at 3000xg for 10 min. The supernatant from each homogenate was divided into two parts, for MDA and protein concentrations, respectively. The protein concentration was determined using the

method described by Lowry *et al.* (1951). For the determination of MDA concentrations, 0.25ml of supernatant was mixed with 0.5ml of 10% trichloroacetic acid, and then heated in a boiling water bath for 15 min. After cooling under running tap water for 5 min, the mixture was centrifuged at 1600xg for 10 min, 1ml of the supernatant was then added to 0.5ml of 6.7g/L TBA solution in a test tube and placed in a boiling water bath for 15 min. The solution was then cooled under running tap water and the absorbance was then measured at 532nm using a UV spectrophotometer (T80<sup>+</sup>UV/VIS Spectrometer<sup>®</sup> PG Instruments Ltd, UK). The MDA concentration was calculated by the absorbance co-efficient, MDA-TBA complex 1.56x10<sup>5</sup>/cm and expressed in nmol/mg of protein.


*Statistical analysis*

Values were expressed as mean±SEM and subjected to one-way analysis of variance followed by Tukey's test. Value of P<0.05 was considered significant.

**Results**

*Effect of treatments on serum glucose concentration*

The effect of treatments on serum glucose is shown in Figure 1. There was no significant difference (P>0.05) between the groups. However, a comparative increase in the concentration of serum glucose was recorded in CPF+ Pb group.



**Figure 1: The effect of corn oil (C/oil), vitamin C (VC) and/or chlorpyrifos (CPF) and lead on serum glucose concentration.**

*Effect of treatments on hepatic glycogen concentration*

The effect of treatments on hepatic glycogen concentration is shown in Figure 2. A significant

(P<0.05) decrease in the hepatic glycogen concentration was recorded in the CPF + Pb group compared to the other group.




Figure 2: Effect of corn oil (C/oil), vitamin C (VC) and/or chlorpyrifos (CPF) + lead (Pb) on hepatic glycogen concentration. <sup>a</sup>P<0.01 vs VC group; <sup>b</sup>P<0.05 vs corn oil; <sup>c</sup>P<0.05 vs VC group.

*Effect of treatments on pancreatic malondialdehyde concentration*

The effect of treatments on pancreatic malondialdehyde concentration is shown in Figure 3. There was a significant increase (P<0.01) in the concentration of pancreatic MDA

in the CPF+Pb group compared to the other groups. Similarly, there was a significant increase (P<0.05) in the MDA concentration in the VC+CPF+Pb group compared to the VC group.




Figure 3: Effect of corn oil (C/oil), vitamin C (VC) and/or chlorpyrifos (CPF) + lead (Pb) on pancreatic malonaldehyde concentration. <sup>ab</sup>P<0.01 vs corn oil and VC groups, respectively; <sup>c</sup>P<0.05 vs VC+CPF+Pb group; <sup>d</sup>P<0.05 vs VC group.

## Discussion

A high glucose concentration was recorded in the group co-administered with CPF and Pb. Hyperglycemia has been recorded following CPF (Ambali, 2009) and lead poisoning (Krishna and Ramachandran, 2009). The hyperglycemic response recorded in the group co-administered with CPF and Pb in the present study may be due to increased pancreatic lipoperoxidation, which ultimately affect the elaboration of insulin that is critical in the regulation of glucose concentration in the blood. Similarly, OP compounds have been shown to increase glycogenolysis by inhibiting insulin activity and stimulating glucagon activity (Rahimi and Abdollahi, 2007). In addition, the elevated serum glucose concentration increases the peroxide levels in the islets, which normally have poor antioxidant enzymes mRNA contents and activity levels (Vousough-Ghanbari *et al.*, 2007). This glucotoxic condition causes a decrease in the levels of two critical regulatory proteins, PDX-1 and MafA (a family of Maf family of transcription factors) that normally bind to the insulin promoter and stimulate insulin gene transcription (Vousough-Ghanbari *et al.*, 2007). Decrease in the levels of these two proteins causes decrease in insulin promoter activity, insulin gene expression and insulin secretion (Harmon *et al.*, 1999; Robertson *et al.*, 2003). This may have contributed to the apparent increase in glucose concentration in CPF+Pb group.

Similarly, the induction of toxic stress may have played some role in the hyperglycemia recorded in the group co-administered CPF and Pb. Stress generally activates the hypothalamo-pituitary-adrenal (HPA) axis and the sympathetic nervous system resulting in hyperglycemia (Bateman *et al.*, 1989; Blalock, 2002; Mechanick, 2006). The activation of HPA axis results in increased elaboration of glucocorticoid from the adrenal cortex, which eventually results in increased glucose concentration through increased gluconeogenesis (Rahimi and Abdollahi, 2007) and impairment of glucose uptake in skeletal muscle (Oda *et al.*, 1995). Similarly, the stimulation of sympathetic nervous system during stress leads to enhanced release of catecholamines, glucagon, and growth hormone which result in promotion of gluconeogenesis, glycogenolysis, insulin resistance, and constitution of hyperglycemia (Gustavson *et al.*, 2003; Gearhart *et al.*, 2006). It has been shown that OP compounds induces insulin resistance

by inhibiting glucose transport in skeletal muscle via impinging on the component of insulin signaling pathway (Chiasson *et al.*, 1981; Hunt and Ivy, 2002). The oxidative stress induced by both CPF and Pb in the body may have been exacerbated by the hyperglycemia through the formation of advanced glycation end products (Rahimi *et al.*, 2005; Gillery, 2006). Therefore, hyperglycemia is considered as a mechanism for development of oxidative stress in OP poisoning (Rahimi and Abdollahi, 2007).

Furthermore, increased muscular activity resulting from increased cholinergic activity (Shi and Screming, 1992; Garg *et al.*, 2004) and pancreatitis due cholinergic stimulations (Harputluoglu *et al.*, 2003; Markrides *et al.*, 2005) have been shown to escalate OP-induced hyperglycemia. Pancreatic  $\beta$ -cells contain muscarinic ACh receptors, which are involved in the glucose-dependent production of insulin (Duttaroy *et al.*, 2004). The cholinergic system plays an essential role in insulin release (Balkan and Dunning, 1995; Ahren *et al.*, 1999; D'Alessio *et al.*, 2001). CPF and lead are known inhibitor of AChE activity (Ambali *et al.*, 2010; Ademuyiwa *et al.*, 2007), the enzyme responsible for the degradation of acetylcholine (ACh). Thus, exposure to the CPF and Pb may have resulted in increased accumulation of ACh, potentially leading to overstimulation and eventual downregulation of its receptors (van Koppen and Kaiser, 2003) and consequent reduction of insulin production (Montgomery *et al.*, 2008). Furthermore, prolonged stimulation by ACh may reduce  $\beta$ -cell sensitivity to glucose (Gilon and Henquin, 2001). AChE inhibition has been shown to be partly responsible for OP-induced hyperglycemia (Pourkahlili *et al.*, 2009; Joshi and Rashini, 2010). Lead has equally been shown to increase glucose synthesis as well as suppressed pancreatic function (Stevenson *et al.*, 1976). All these factors either individually or collectively may have contributed to increased glucose concentration in group co-administered CPF and Pb.

Pretreatment with vitamin C restored the serum glucose concentration to apparently normal level, indicating the role of oxidative stress in the hyperglycemic response observed in rats exposed to a combination of CPF and Pb. This may be due to the ability of the antioxidant vitamin to prevent oxidative damage to the pancreatic islets. A previous study has shown that  $\alpha$ -tocopherol, a lipid soluble antioxidant vitamin restored diazinon-induced insulin

secretion (Pourkhalili *et al.*, 2009). Apart from lipoperoxidation, vitamin C has been shown to restore AChE activity inhibited by OPs (Yavuz *et al.*, 2004; Ambali *et al.*, 2010d). This may have apparently restored the level of ACh, a potent secretagogue of both insulin and glucagon (Duttaroy *et al.*, 2004). This may have aided pancreatic  $\beta$  cell activity, increasing the elaboration of insulin and therefore suppressing the hyperglycemia. Some other non-antioxidant activity of vitamin C may have complemented the restoration of normal serum glucose level following alteration by CPF and Pb.

The present study has also revealed a decrease in hepatic glycogen concentration in the group co-administered CPF and Pb. This may be due to pancreatic damage as a result of lipoperoxidative changes, which causes an increase in blood glucose concentration and a decrease in intracellular glucose that can be used to synthesize glycogen. Furthermore, studies have shown that CPF causes impairment in hepatic function due to oxidative changes (Goel *et al.*, 2006; Ambali *et al.*, 2007; Ambali, 2009). This reduces the ability of the liver to synthesize glycogen. OP compounds also causes inhibition of AChE activity at the neuroeffector sites in the adrenal medulla, resulting in increased adrenalin secretion (Gupta, 1974) and consequent elevation in glycogenolytic processes in the liver and skeletal muscle (Gustavson *et al.*, 2003). Similarly, OP compounds have been shown to increase lipolysis, resulting in the elevation of free fatty acids that is known to have an inhibitory effect on the insulin signaling and inhibit glycogen synthesis (Itani *et al.*, 2002). In addition, oxidative stress has been shown to induce an impairment of insulin action in glucose transport and glycogen synthesis. The decrease in insulin-stimulated glycogen synthesis during oxidative stress has been linked to the impairment of insulin to stimulate the activity of glycogen synthase (Dokken *et al.*, 2008). The consequences of these are hyperglycemia and reduced glycogen synthesis.

Pretreatment with vitamin C restored the hepatic glycogen level to apparently normal level, indicating the role of oxidative stress in the depleted hepatic glycogen reserves observed in rats exposed to combination of CPF and Pb. The cytoprotection offered by vitamin C pretreatment on oxidative stress-induced hepatic and pancreatic damage may have played a very significant role in the apparent restoration of hepatic glycogen reserves in the vitamin C

pretreated group. In addition, the ability of vitamin C to restore AChE activity (Yavuz *et al.*, 2004; Ambali *et al.*, 2010d) may have reduced the activity of the adrenal medulla to elaborate adrenalin and consequently reduced hepatic glycogenolysis.

The increase in pancreatic MDA concentration in the group co-administered CPF and Pb was an indication of lipoperoxidation. Oxidative stress is being increasingly implicated in the pathogenesis of pancreatic inflammation (Leung and Chan, 2009). The lipoperoxidative damage to the pancreas may have compromised its structural and functional integrity. Diazinon, an OP insecticidal compound, had equally been shown to reduce glucose-stimulated insulin secretion through the induction of oxidative and toxic stress in the islet of Langerhans (Pourkhalili *et al.*, 2009). In addition to being a potential inhibitor of insulin gene transcription and suppressor of the promoter, reactive oxygen species (ROS) also causes a decrease in mRNA and protein, and impairs the homeodomain transcription factor pancreatic/duodenal homeobox-1 (PDX-1) activity (Matsuoka *et al.*, 1997), which is an important insulin transcription factor. The result is a decreased insulin secretion and consequent hyperglycemia.

Pretreatment with vitamin C restored the pancreatic MDA concentration to apparently normal level, affirming the role of oxidative stress in the pancreatic lipoperoxidative response observed in rats exposed to combinations of CPF and Pb. Vitamin C prevented oxidative damage to the pancreatic islets, thereby aiding in the maintenance of its structural and functional status. Vitamin C, like many other nutritional antioxidants act as free radical scavengers by directly neutralizing them, reduce peroxide concentration and repair oxide membranes and by quenching iron to decrease ROS production. Furthermore, they are known to regulate a number of genes and signal regulatory pathways thereby preventing the incidence of cell death (Young and Woodside, 2001).

### Conclusion

The present study has shown that vitamin C pretreatment restored the hyperglycemia and decreased liver glycogen reserve induced by subchronic co-administration of CPF and Pb. This may be partly due to antioxidant property of the vitamin, which prevented oxidative damage to the pancreas. This study therefore

underscores the role of oxidative stress in the etiopathogenesis of hyperglycemia following exposure to combination of CPF and Pb.

## References

Ademuyiwa O, Ugbaja RN, Rotimi SO, Abam E, Okediran BS, Dosumu OA, Onunkwor BO, 2007. Erythrocyte acetylcholinesterase activity as a surrogate indicator of lead-induced neurotoxicity in occupational lead exposure in Abeokuta, Nigeria. *Environmental Toxicology and Pharmacology*, 24(2): 183-188.

Ahren B, Sauerberg P, Thomsen C, 1999. Increased insulin secretion and normalization of glucose tolerance by cholinergic agonism in high fat-fed mice. *American Journal of Physiology*, 277: E93-102.

Ambali SF, 2009. Ameliorative effect of antioxidant vitamins C and E on neurotoxicological, haematological and biochemical changes induced by chronic chlorpyrifos exposure in Wistar rats. PhD Dissertation, Ahmadu Bello University, Zaria, Nigeria.

Ambali SF, Abubakar AT, Shittu M, Yaqub LS, Anafi SB, Abdullahi A, 2010a. Chlorpyrifos-induced alteration of hematological parameters in Wistar rats: Ameliorative effect of zinc. *Research Journal of Environmental Toxicology*, 4(2): 55-66.

Ambali SF, Abubakar AT, Shittu M, Yaqub LS, Kobo PI, Giwa A, 2010b. Zinc ameliorates chlorpyrifos-induced erythrocyte fragility and lipoperoxidative changes in Wistar rats. *New York Science Journal*, 3: 117-122.

Ambali S, Akanbi D, Igbokwe N, Shittu M, Kawu M, Ayo J, 2007. Evaluation of subchronic chlorpyrifos poisoning on haematological and serum biochemical changes in mice and protective effect of vitamin C. *The Journal of Toxicological Science*, 32(2): 111-120.

Ambali SF, Ayo JO, Ojo SA, Esievo KAN, 2010c. Ameliorative effect of vitamin C on chlorpyrifos-induced increased erythrocyte fragility in Wistar rats. *Human and Experimental Toxicology*. Epub ahead of print

Ambali SF, Idris SB, Onukak C, Shittu M, Ayo JO, 2010d. Ameliorative effects of vitamin C on short-term sensorimotor and cognitive changes induced by acute chlorpyrifos exposure in Wistar rats. *Toxicology and Industrial Health*, 26(9): 547-558.

Balkan B, Dunning BE, 1995. Muscarinic stimulation maintains *in vivo* insulin secretion in response to glucose after prolonged hyperglycemia. *American Journal of Physiology*, 268: R475-479.

Bateman A, Singh A, Kral T, Solomon S, 1989. The immune-hypothalamic-pituitary-adrenal axis. *Endocrinology Review*, 10: 92-112.

Blalock JE, 2002. Harnessing a neural-immune circuit to control inflammation and shock. *Journal of Experimental Medicine*, 195: F25-F28.

Casida JE, Quistad GB, 2004. Organophosphate toxicity: safety aspects on non acetylcholinesterase secondary targets. *Chemical Research in Toxicology*, 17: 983-998.

Chiasson JL, Shikama H, Chu DT, Exton JH, 1981. Inhibitory effect of epinephrine insulin-stimulated glucose uptake by rat skeletal muscle. *Journal of Clinical Investigations*, 68: 706-713.

D'Alessio DA, Kieffer TJ, Taborsky GJ, Havel PJ, 2001. Activation of the parasympathetic nervous system is necessary for normal meal-induced insulin secretion in rhesus macaques. *Journal of Clinical Endocrinology and Metabolism*, 86: 1253-1259.

Dokken BB, Saengsirisuwan V, Kim JS, Teachey MK, Henrikson EJ, 2008. Oxidative stress-induced insulin resistance in rat skeletal muscle: role of glycogen synthase kinase-3. *American Journal of Physiology, Endocrinology and Metabolism*, 294: 615-621.

Draper HH, Hadley M, 1990. Malondialdehyde determination as index of lipid peroxidation. *Methods in Enzymology*, 186: 421-431.

Duttaray A, Zimiliki CL, Gautam D, Cui Y, Mears D, Wess J, 2004. Muscarinic stimulation of pancreatic insulin and glucagon release is abolished in M3 muscarinic acetylcholine receptor deficient mice. *Diabetes*, 53: 1714-1720.

Ercal N, Treratphan P, Hammond TC, Mathews RH, Grannemann NH, Spitz DR, 1996. *In vivo* indices of oxidative stress in lead exposed C57BL/6 mice are reduced by treatment with meso-2, 3-dimercaptosuccinic acid or N-acetyl cysteine. *Free Radical Biology and Medicine*, 21: 157-161.

Garcia SJ, Seidler FJ, Slotkin TA, 2003. Developmental neurotoxicity elicited by prenatal or postnatal chlorpyrifos exposure: effects on neurospecific proteins indicate changing vulnerabilities. *Environmental Health Perspectives*, 111: 297-303.

Garg UK, Pal AK, Jha GJ, Jadhao SB, 2004. Haemato-biochemical and immunopathophysiological effects of chronic toxicity with synthetic pyrethroid, organophosphate, and chlorinated pesticides in broiler chicks. *International Immunopharmacology*, 4: 1709-1722.

Gearhart MM, Parbhoo SK, 2006. Hyperglycemia in the critically ill patient, AACN. Clinical Issues, 17: 50–55.

Gillery P, 2006. Oxidative stress and protein glycation in diabetes mellitus. Annales de Biologie Clinique (Paris), 64: 309–314.

Gilon P, Henquin JC, 2001. Mechanisms and physiological significance of the cholinergic control of pancreatic beta-cell function. Endocrine Review, 22: 565–604.

Goel A, Danni V, Dhawan DK, 2006. Role of zinc in mitigating the toxic effects of chlorpyrifos on hematological alterations and electron microscopic observations in rat blood. BioMetals, 19(5): 483-492.

Good CA, Krames H, Somongi M, 1933. Chemical Procedure for Analysis of Polysaccharides. Methods in Enzymology, VII 34.

Gultekin F, Delibas N, Yasar S, Kilinc I, 2001. *In vivo* changes in antioxidant systems and protective role of melatonin and a combination of vitamin C and vitamin E on oxidative damage in erythrocytes induced by chlorpyrifos-ethyl in rats. Archives of Toxicology, 75(2): 88-96.

Gupta PK, 1974. Malathion induced biochemical changes in rats. Acta Pharmacology and Toxicology, 35: 191-194.

Gustavson SM, Chu CA, Nishizawa M, Farmer B, Neal D, Yang Y, Donahue EP, Flakoll P, Cherrington AD, 2003. Interaction of glucagon and epinephrine in the control of hepatic glucose production in the conscious dog. American Journal of Physiology, Endocrinology and Metabolism, 284: E695–E707.

Harmon JS, Gleason CE, Tanaka Y, Oseid EA, Hunter-Berger KK, Robertson RP, 1999. *In vivo* prevention of hyperglycemia also prevents glucotoxic effects on PDX-1 and insulin gene expression. Diabetes, 48: 1995–2000.

Harputluoglu MM, Kantarceken B, Karıncaoglu M, Aladag M, Yildiz R, Ates M, Yildirim B, Hilmioglu F, 2003. Acute pancreatitis: an obscure complication of organophosphate intoxication. Human and Experimental Toxicology, 22: 341–343.

Hayes MM, van der Westhuizen NG, Gelfand M, 1978. Organophosphate poisoning in RhoOdesia. A study of the clinical features and management of 105 patients. South Africa Medical Journal, 54: 230-234.

Houston DK, Johnson MA, 2000. Does vitamin C intake protect against lead toxicity? Nutrition Review, 58(3): 73-75.

Hunt DG, Ivy JL, 2002. Epinephrine inhibits insulin-stimulated muscle glucose transport. Journal of Applied Physiology, 93: 1638–1643.

Itani SB, Ruderman, NB, Schmieder F, Boden G, 2002. Lipid-induced insulin resistance in human muscle is associated with changes in diacylglycerol, protein kinase C, and I $\kappa$ B $\alpha$ . Diabetes, 51: 2005–2011.

Joshi AKR, Rajini PS, 2010. Hyperglycemic and stressogenic effects of monocrotophos in rats: Evidence for the involvement of acetylcholinesterase inhibition. Experimental and Toxicologic Pathology, Article in Press.

Krishna H, Ramachandran AV, 2009. Biochemical alterations induced by the acute exposure to combination of chlorpyrifos and lead in Wistar rats. Biology and Medicine, 1(2): 1-6.

Lowry H, Rosebrough NJ, Farr AL, Randall RJ, 1951. Protein measurements with the folin phenol reagent. Journal of Biological Chemistry, 193: 265–275.

Leung SP, Chan YC, 2009. Role of oxidative stress in pancreatic inflammation. Antioxidants & Redox Signaling, 11(1): 135-165.

Makrides C, Koukouvas M, Achillews G, Tsikkos S, Vounou E, Symeonides M, Christodoulides P, Ioannides M, 2005. Methomyl-induced severe acute pancreatitis: possible etiological association. Journal of the Pancreas, 6: 166–171.

Matsuoka TA, Kajimoto Y, Watada H, Kaneto H, Kishimoto M, Umayahara Y, Fujitani Y, Kamada T, Kawamori R, Yamasaki Y, 1997. Glycation-dependent, reactive oxygen species-mediated suppression of the insulin gene promoter activity in HIT cells. Journal of Clinical Investigations, 99(1):144–150.

Mechanick JI, 2006. Metabolic mechanisms of stress hyperglycemia. Journal of Parenteral and Enteral Nutrition, 30: 157–163.

Meller D, Fraser I, Kryger M, 1981. Hyperglycemia in anticholinergic poisoning. Canadian Medical Association Journal, 124: 745–748.

Montgomery MP, Kamel F, Saldana TM, Alavanja MCS, Sandler DP, 2008. Incident diabetes and pesticide exposure among licensed pesticide applicators: Agricultural Health Study, 1993–2003. American Journal of Epidemiology, 67(10): 1235–1246.

Namba T, Nolte CT, Jackrel J, Grob D, 1971. Poisoning due to organophosphate insecticides.

Acute and chronic manifestations. American Journal of Medicine, 50(4): 475-492.

Oda N, Nakai A, Mokuno T, Sawai Y, Nishida Y, Mano T, Asano K, Itoh Y, Kotake M, Kato S, 1995. Dexamethasone-induced changes in glucose transporter 4 in rat heart muscle, skeletal muscle and adipocytes. European Journal of Endocrinology, 133: 121-126.

Oladipo OO, 2010. Ameliorative effects of ascorbic acid on neurobehavioural, haematological and biochemical changes induced by subchronic lead exposure in Wistar rats. MSc thesis, Ahmadu Bello University, Zaria, Nigeria.

Olaleye SB, Adaramoye OA, Erigbali PP, Adeniyi OS, 2007. Lead exposure increases oxidative stress in the gastric mucosa of HCl/ethanol-exposed rats. World Journal of Gastroenterology, 13(38): 5121-5126.

Pope CN, 1999. Organophosphorus pesticides: Do they all have the same mechanism of toxicity? Journal of Toxicology and Environmental Health B Critical Reviews, 2: 161-181.

Pourkhalili N, Poumourmohammadi S, Rahimi F, Vosough-Ghanbari S, Baeri M, Ostad NS, Abdollahi M, 2009. Comparative effects of calcium channel blockers, autonomic nervous system blockers, and free radical scavengers on diazinon-induced hyposecretion of insulin from isolated islets of Langerhans in rats. Arh Hig Rada Toksikol, 60: 157-164.

Rahimi R, Abdollahi M, 2007. A review on the mechanisms involved in hyperglycaemia induced by organophosphorus pesticide. Pesticide Biochemistry and Physiology, 88: 115-121.

Rahimi R, Nikfar S, Larijani B, Abdollahi M, 2005. A review on the role of antioxidants in the management of diabetes and its complications. Biomedicine and Pharmacotherapy, 59: 365-373.

Robertson RP, Harmon J, Tran PO, Tanaka Y, Takahashi H, 2003. Glucose toxicity in beta cells: type 2 diabetes, good radicals gone bad, and the glutathione connection. Diabetes, 52: 581-587.

Seifert J, 2001. Toxicologic significance of the hyperglycemia caused by organophosphorus insecticides. Bulletin of Environmental Contamination and Toxicology, 67: 463-469.

Shaffi SA, 1979. Lead toxicity, biochemical and physiological imbalance in nine fresh water teleosts. Toxicology Letters, 4: 155-161.

Shih TM, Scremin OU, 1992. Cerebral blood flow and metabolism in soman-induced convulsions. Brain Research Bulletin, 28: 735-742.

Stevenson A, Merali Z, Kacew S, Singhal RL, 1976. Effects of subacute and chronic lead treatment on glucose homeostasis and renal cyclic AMP metabolism in rats. Toxicology, 6(3): 265-275.

Van Koppen CJ, Kaiser B, 2003. Regulation of muscarinic acetylcholine receptor signaling. Pharmacology and Therapeutics, 98: 197-220.

Vosough-Ghanbari S, Sayyar P, Poumourmohammadi S, Aliahmadi S, Ostad NS, Abdollahi M, 2007. Stimulation of insulin and glucagon synthesis in rat Langerhans islets by malathion *in vitro*: Evidence for mitochondrial interaction and involvement of subcellular non-cholinergic mechanisms. Pesticide Biochemistry and Physiology, 89: 130-136.

Vojarova de Courten B, Weyer C, Stefan N, Horton M, Delparigi A, Havel P, Bogardus C, Tataranni PA, 2004. Parasympathetic blockade attenuates augmented pancreatic polypeptide but not insulin secretion in Pima Indians. Diabetes, 53: 663-671.

Wang J, Junhui W, Zhaoming Z, 2006. Oxidative stress in mouse brain exposed to lead. Annals of Occupational Hygiene, 50(4): 405-409.

Yavuz T, Delibao N, Yâldâr B, Altuntao I, Candâr O, Cora A, Karahan N, Âbrioim E, Kutsal A, 2004. Vascular wall damage in rats induced by methidathion and ameliorating effect of vitamins E and C. Archives of Toxicology, 78: 655-659.

Young IS, Woodside IS, 2001. Antioxidant in health and disease. Journal of Clinical Pathology, 54: 176-186.